

From the office of Drs. Brickner, Menning, Moeller, Coffey and Julien:

If you are on any blood thinners, e.g. PLAVIX, LOVENOX, TICLID, COUMADIN, please be certain you have notified the office. Stop medication as directed.

If you are taking any of the medications listed below, they would need to be stopped prior to any scheduled procedure either 1 or 2 weeks before.

MEDICATIONS TO STOP ONE (1) WEEK BEFORE

ALL over-the-counter medications (EXCEPT TYLENOL products). *This can include any of the following (and some that may not be listed):*

ASPIRIN
EXCEDRIN
IRON PILLS
VITAMIN E (Alone)
MOTRIN
ADVIL
ALEVE
IBUPROFEN

Stop NON-STEROIDAL / ANTI-INFLAMMATORY medications such as:

DAYPRO
VOLTAREN
FELDENE
CLINORIL
NAPROSYN
RELAFEN

HERBALS & MEDICATIONS TO BE OFF TWO (2) WEEKS

PRIOR TO GENERAL ANESTHESIA, PER DEPARTMENT OF ANESTHESIA

MEDICATIONS	HERBALS
1. DIET PILLS	1. GINSENG
2. FENFLURAMINE	2. FEVERFEW
3. PHENTERMINE	3. GINGER
4. EPHEDRA	4. GOLDSEAL
5. PONDIMIN	5. LICORICE
6. IONAMIN	6. VALERIAN
7. FASTIN	7. GARLIC
8. METABOLIFE	8. KAVA KAVA
9. REDUX	9. ST. JOHNS WORT
10. CENTRUM PERFORMANCE	10. GINKO BILOBA
11. MERIDIA	